



Proud supplier of:

Spinosad

Topical Suspension 0.9%



**One product.
Two indications.¹
No reported resistance.^{2,3}**

Safe and effective treatment for head lice and scabies.^{1,2}

Product ordering information for 120 mL (4 oz.) bottles

Dosing	Strength	Pack	NDC	Item No.
One Bottle	0.9%	120 mL Bottle	52246-570-04	5303797



**Spinosad Topical Suspension
Prescribing Information**

cipher™
PHARMACEUTICALS

See back page for Important Safety Information

HEAD LICE

IMPORTANT SAFETY INFORMATION

INDICATION

Spinosad Topical Suspension 0.9% is a pediculicide indicated for the topical treatment of head lice infestations in adult and pediatric patients six (6) months of age and older.

ADJUNCTIVE MEASURES

Spinosad Topical Suspension 0.9% should be used in the context of an overall lice management program:

- Wash in hot water or dry-clean all recently worn clothing, hats, used bedding and towels
- Wash personal care items such as combs, brushes, and hair clips in hot water
- A fine-tooth comb or special nit comb may be used to remove dead lice and nits

IMPORTANT SAFETY INFORMATION

Spinosad Topical Suspension 0.9% contains benzyl alcohol and is not recommended for use in neonates and infants below the age of 6 months. Intravenous administration of benzyl alcohol has been associated with serious adverse reactions and death in neonates and low birth-weight infants. Most common adverse events were: application site redness (3%), eye redness (2%) and application site irritation (1%).

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

**Application
instructions for
head lice**



SCABIES

IMPORTANT SAFETY INFORMATION

INDICATION

Spinosad Topical Suspension 0.9% is a scabicide indicated for the topical treatment of scabies infestations in adult and pediatric patients four (4) years of age and older.

ADJUNCTIVE MEASURES

- Wash in hot water or dry-clean any bedding, clothing and towels used by anyone having scabies.
- If any member of a household presents with scabies, all household members should be treated with Spinosad Topical Suspension 0.9%.

IMPORTANT SAFETY INFORMATION

Spinosad Topical Suspension 0.9% contains benzyl alcohol and is not recommended for use in pediatric patients below the age of 4 years. The safety and effectiveness of Spinosad Topical Suspension 0.9% have not been established in pediatric patients less than 4 years of age with scabies infestation. Most common adverse events were: application site irritation (3%)* and dry skin (2%).

*Application site irritation also includes application site pain and burning sensation.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

**Application
instructions for
scabies**



References: 1. Spinosad Prescribing Information. 2. Spinosad at 0.9% in the treatment of scabies: Efficacy results from 2 multicenter, randomized, double-blind, vehicle-controlled studies, Seiler, Jeffrey C. et al. *Journal of the American Academy of Dermatology*, Volume 86, Issue 1, 97 – 103, 2022 3. Spinosad Technical Bulletin Dow AgroSciences LLC (Now Corteva agri-science), Form No. Y45-000-001 (01/01) CBK. 7.

Spinosad
Topical Suspension 0.9%

cipher[™]
PHARMACEUTICALS

Manufactured and Distributed by Cipher Pharmaceuticals.
© 2025 Cipher Pharmaceuticals US LLC. All Rights Reserved
SPN-WVDC-000



INDEPENDENT
PHARMACY
COOPERATIVE

Proud supplier of:

Natroba[™]
(spinosad) Topical Suspension, 0.9%



**One product.
Two indications.¹
No reported resistance.^{2,3}**

Safe and effective treatment for head lice and scabies.^{1,2}

Product ordering information for 120 mL (4 oz.) bottles

Dosing	Strength	Pack	NDC	Item No.
One Bottle	0.9%	120 mL Bottle	52246-0929-04	5303748



Natroba
Prescribing
Information

cipher[™]
PHARMACEUTICALS

See back cover for Important Safety Information

HEAD LICE

IMPORTANT SAFETY INFORMATION

INDICATION

Natroba™ Topical Suspension is a pediculicide indicated for the topical treatment of head lice infestations in adult and pediatric patients six (6) months of age and older.

ADJUNCTIVE MEASURES

Natroba™ Topical Suspension should be used in the context of an overall lice management program:

- Wash in hot water or dry-clean all recently worn clothing, hats, used bedding and towels
- Wash personal care items such as combs, brushes, and hair clips in hot water
- A fine-tooth comb or special nit comb may be used to remove dead lice and nits

IMPORTANT SAFETY INFORMATION

Natroba™ Topical Suspension contains benzyl alcohol and is not recommended for use in neonates and infants below the age of 6 months. Intravenous administration of benzyl alcohol has been associated with serious adverse reactions and death in neonates and low birth-weight infants. Most common adverse events were: application site redness (3%), eye redness (2%) and application site irritation (1%).

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Application
instructions for
head lice



SCABIES

IMPORTANT SAFETY INFORMATION

INDICATION

Natroba™ Topical Suspension is a scabicide indicated for the topical treatment of scabies infestations in adult and pediatric patients four (4) years of age and older.

ADJUNCTIVE MEASURES

- Wash in hot water or dry-clean any bedding, clothing and towels used by anyone having scabies.
- If any member of a household presents with scabies, all household members should be treated with Natroba™ Topical Suspension.

IMPORTANT SAFETY INFORMATION

Natroba™ Topical Suspension contains benzyl alcohol and is not recommended for use in pediatric patients below the age of 4 years. The safety and effectiveness of Natroba™ Topical Suspension have not been established in pediatric patients less than 4 years of age with scabies infestation. Most common adverse events were: application site irritation (3%)* and dry skin (2%).

*Application site irritation also includes application site pain and burning sensation.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Application
instructions for
scabies



References: 1. Natroba Prescribing Information. 2. Spinosad at 0.9% in the treatment of scabies: Efficacy results from 2 multicenter, randomized, double-blind, vehicle-controlled studies. Seiler, Jeffrey C. et al. *Journal of the American Academy of Dermatology*, Volume 86, Issue 1, 97 – 103, 2022 3. Spinosad Technical Bulletin Dow AgroSciences LLC (Now Corteva agriscience), Form No. Y45-000-001 (01/01) CBK. 7.

Natroba™
(spinosad) Topical Suspension, 0.9%

cipher™
PHARMACEUTICALS

Manufactured and Distributed by Cipher Pharmaceuticals.
© 2025 Cipher Pharmaceuticals US LLC. All Rights Reserved
NAT-WVDC-000